

Listing of Claims

1. (currently amended) A method for selectively labeling or tagging one or more phosphate groups in a natural or synthetic peptide or protein ~~in the presence of which also contains~~ one or more carboxylic acid groups or esters thereof which method comprises the steps of:

- a. reacting the natural or synthetic peptide or protein with a protective group that is an amine that reacts to protect the one or more phosphate groups therein by forming phosphoramidate bonds and to protect any carboxylic acid groups and esters thereof therein by forming amide bonds;
- b. treating the protected peptide or protein under conditions which selectively substantially cleave the phosphoramidate ~~bond~~ bonds, without substantially cleaving the amide ~~bond~~ bonds, to regenerate one or more free phosphate groups in the peptide or protein; and
- c. reacting the one or more free phosphate groups in the peptide or protein, in which the carboxylic acid groups remain protected, with a label or tag comprising a functional group that reacts with a phosphate group to thereby generate a peptide or protein in which the phosphate groups are selectively labeled or tagged.

2. canceled

3. (currently amended) The method of claim 1 wherein the label or tag is a solid phase material and ~~a free phosphate group~~ the one or more free phosphate groups of the ~~oligomer or polymer is~~ peptide or protein are covalently linked to the solid phase material directly or indirectly through a linker moiety.

4. (currently amended) The method of claim 1 wherein the amino group of the protective group that is an amine is reacted with the one or more phosphate group groups of the peptide or protein using a carbodiimide catalyzed reaction.
5. (currently amended) The method of claim 3 ~~claim 1~~ wherein the one or more free phosphate groups of the peptide or protein are reacted with a linker group ~~having wherein~~ the linker group comprises a phosphate reactive group and a second reactive group that functions for forming a covalent bond to the solid phase material.
6. (original) The method of claim 5 wherein the second reactive group is a sulfhydryl reactive group.
7. (currently amended) The method of claim 1 wherein the ~~amine~~-protecting group is ethanolamine.
8. (currently amended) The method of claim 7 wherein in step b the protected peptide or protein is treated with trifluoroacetic acid to selectively regenerate one or more free phosphate groups.
9. (currently amended) The method of claim 1 wherein the one or more free phosphate groups are reacted with a linker ~~containing~~ which contains a sulfhydryl group or which contains a latent reactive group that can be transformed into a sulfhydryl group.
10. (currently amended) The method of claim 9 wherein the one or more free phosphate group is-groups are first reacted with cystamine and thereafter a free one or more free sulfhydryl group is-groups are generated by reduction of the disulfide group of the cystamine.
11. (original) The method of claim 10 wherein the reducing agent is DTT.

12. (currently amended)The method of claim 9 wherein in step c the peptide or protein is covalently attached to ~~the~~ a solid support material through reaction with the sulfhydryl group of the linker.
13. (original)The method of claim 12 wherein the solid support material is glass beads with immobilized iodoacetyl groups.
14. (currently amended) The method of claim 1 wherein the natural or synthetic peptide or protein is ~~peptides or proteins are~~ obtained from a tryptic digest.
15. (original)The method of claim 1 wherein the label or tag is a radiolabel, a fluorescent label, a colorimetric label or an affinity label.
16. (previously presented) The method of claim 1 wherein the label or tag is an affinity label.
17. (original)The method of claim 1 wherein the label is a reactive label.
18. (currently amended)The method of claim 1 further comprising a step of detecting the selectively labeled ~~phosphopeptide or~~ tagged peptide or protein.
19. (currently amended)The method of claim 18 wherein the ~~phosphopeptide~~ selectively labeled or tagged peptide or protein is detected by detection of the label or tag.
20. (currently amended)The method of claim 18 wherein the label or tag is an affinity label and the selectively labeled or tagged peptide or protein ~~phosphopeptide~~ is detected by binding to a ~~corresponding~~ capture reagent.
21. (currently amended)The method of claim 1 ~~further comprising a step of separating~~

~~selectively labeled phosphopeptides from the mixture by attachment of the phosphopeptide- wherein step c comprises attaching the-peptide or protein having one or more free phosphate groups to a solid support or by-binding of the-phosphopeptide the peptide or protein having one or more free phosphate groups to a capture reagent.~~

22. (currently amended)A method for detecting one or more phosphopeptides or phosphoproteins in one or more samples containing a mixture of peptides, proteins or both which comprises the steps of:
- a. selectively protecting the carboxylic acid groups or esters thereof of the peptides or proteins in the one or more samples such that any phosphate groups in the peptides or proteins are ~~remain~~ unprotected;
 - b. selectively labeling the unprotected phosphate groups in the peptides or proteins in the sample with a label having a functional group that reacts directly or indirectly with a phosphate group; and
 - c. detecting the peptides or proteins carrying the selective label of step b to detect the one or more phosphopeptides or phosphoproteins in the sample.
23. (currently amended)The method of claim 22 wherein, in step a, the carboxylic acid groups or esters thereof of the peptides or proteins are protected by initial reaction with a protecting group that protects both carboxylic acid groups, esters thereof and phosphate groups in the peptides and proteins followed by selective deprotection of the phosphate groups in the peptides and proteins.
24. (currently amended)The method of claim 23 wherein the carboxylic acid groups, esters thereof and phosphate groups of the peptides or proteins are protected with an amine group that forms amide bonds with carboxylic acid groups and esters thereof and

phosphoramidate bonds with phosphate groups and wherein the protected phosphate groups are thereafter selectively deprotected by cleavage of the phosphoramidate bonds.

25. (currently amended) The method of claim 24 wherein the amide bonds and phosphoramidate bonds are formed by a carbodiimide-catalyzed condensation reactions.
26. (currently amended) The method of claim 25 wherein the phosphate groups are selectively deprotected without cleavage of the amide bonds by treatment with mild acid.
27. (original) The method of claim 22 wherein the label is a radiolabel, a fluorescent label, a colorimetric label or an affinity label.
28. (currently amended) The method of claim 22 wherein the label is an affinity label and the selectively labeled phosphopeptide is phosphopeptides or phosphoproteins are detected by binding to a corresponding capture reagent.
29. (original) The method of claim 22 wherein the label is a reactive label.
30. (original) The method of claim 29 wherein the reactive label carries a reactive group that can form a covalent bond to a solid phase material.
31. (original) The method of claim 29 wherein the reactive label carries a latent reactive group.
32. (currently amended) The method of claim 22 further comprising the step of separating selectively labeled peptides-phosphopeptides or phosphoproteins from the mixture of peptides or proteins in a sample prior to detection step c.
33. (currently amended) The method of claim ~~22~~ 32 wherein the label carries a reactive group

that can form a covalent bond to a solid phase material and wherein the selectively labeled ~~peptides~~ phosphopeptides or phosphoproteins are separated by first covalently attaching the labeled ~~peptides~~ phosphopeptides or phosphoproteins to the solid phase material, then washing the solid support to remove peptides or proteins that are not covalently attached to the support and thereafter releasing ~~peptides covalently attached~~ phosphopeptides or phosphoproteins from the solid phase material.

34. (currently amended)The method of claim 33 wherein the ~~peptides~~ phosphopeptides or phosphoproteins released from the solid phase material are detected using mass spectrometric techniques.
35. (currently amended)The method of claim 34 wherein tandem mass spectrometry is used to detect the ~~peptides~~ phosphopeptides or phosphoproteins.
36. (currently amended)The method of claim 35 wherein tandem mass spectrometry is further used to determine the amino acid sequence of ~~peptides~~ phosphopeptides or phosphoproteins and the precise position of the phosphorylated amino acid within the ~~peptide~~ sequence of the phosphopeptides or phosphoproteins.
37. (currently amended)The method of claim 32 for detecting one or more phosphopeptides or phosphoproteins in two or more samples wherein differentially isotopically labeled ~~carboxylic acid-protecting groups~~ for carboxylic acids or esters thereof are employed with different samples.
38. (currently amended)The method of claim 37 for detecting one or more phosphopeptides or phosphoproteins in two or more samples wherein the labels employed in different samples are differentially isotopically labeled.
39. (currently amended)The method of claim 37 wherein tandem mass spectrometry is used

to detect the one or more phosphopeptides or phosphoproteins and the relative amounts of phosphopeptides or phosphoproteins in the two or more samples is determined by measuring the relative amounts of the differentially isotopically labeled labels present.

40. (currently amended)The method of claim 39 wherein combined microcapillary liquid chromatography and tandem mass spectrometry are employed to detect the ~~peptides~~ phosphopeptides or phosphoproteins.
41. (currently amended)The method of claim 1 wherein the amine-~~group~~ is a hydroxy amine.
42. canceled
43. (currently amended) The method of claim 1 wherein the protecting group ~~comprises a combination of~~ is a differentially isotopically labeled protecting group.
44. (currently amended) The method of claim 43 wherein the protecting group is a hydroxy ~~acid~~-amine.
45. (currently amended)The method of claim 43 wherein the protecting ~~groups are~~ group is differentially isotopically labeled using deuterium.
- C1 46. (original)The method of claim 22 for detecting one or more phosphopeptides in two or more samples wherein differentially isotopically labeled ethanolamine is used to protect the carboxylic acid groups of peptides in different samples.
47. (original)The method of claim 22 further comprising a step of determining the sequence of one or more phosphopeptides detected.

48. (original) The method of claim 47 wherein the sequence of the phosphopeptide is determined by tandem mass spectrometry.
49. (previously presented) The method of claim 48 wherein the samples are protein digests containing peptides and the sequence of the phosphopeptide detected is used to identify the protein from which the phosphopeptide is derived.
50. (currently amended) The method of claim 22 in which the amount of one or more ~~proteins~~ phosphoproteins in a sample is also determined by mass spectrometry, and which further comprises the step of introducing into a sample a known amount of one or more internal standards for each of the ~~proteins~~ phosphoproteins to be quantitated.
51. (original) The method of claim 50 in which different samples represent proteins expressed in response to different environmental or nutritional conditions, different chemical or physical stimuli or at different times.
52. (currently amended) The method of claim 22 wherein phosphopeptides or phosphoproteins in different samples are labeled with different fluorescent labels and the relative ~~intensity~~ amounts of a labeled ~~peptide~~ phosphopeptide or phosphoprotein in different samples can be measured by measuring the relative intensity of the fluorescence emission of ~~the different a labeled peptides~~ phosphopeptide or phosphoprotein in different samples.

C1

Claims 53-67 (canceled)

68. (currently amended) The method of claim 1 further comprising a step of protecting ~~the~~ any amine groups in the peptide or protein prior to step a.

69. (previously presented) The method of claim 68 wherein the amine groups are protected with t-boc or f-moc.
70. (currently amended) The method of claim 22 further comprising a step of protecting the any amine groups in the peptides or proteins of the one or more samples prior to step a.
71. (previously presented) The method of claim 70 wherein the amine groups of the peptides or proteins of the one or more samples are protected with t-boc or f-moc.

Claims 72-73 (canceled)

74. (newly added) The method of claim 21 wherein the peptide or protein having one or more free phosphate groups is present in a mixture containing peptides or proteins that do not contain phosphate groups and which further comprising a step of separating peptides or proteins in which the phosphate groups are selectively labeled or tagged from peptides or proteins that do not contain phosphate groups.
75. (newly added) The method of claim 74 wherein the peptide or protein in which the phosphate groups are selectively labeled or tagged is attached to a solid support and the peptide or protein attached to the solid support is separated from peptides and proteins not attached to the support.
76. (newly added) The method of claim 74 wherein the peptide or protein in which the phosphate groups are selectively labeled or tagged comprises an affinity label which is attached to a capture reagent and the peptide or protein attached to the capture reagent is separated from peptides and proteins not attached to the capture reagent.

77. (newly added) The method of claim 22 for detection of one or more phosphopeptides.
78. (newly added) The method of claim 22 wherein the mixture of peptides and proteins of the one or more samples is obtained from a tryptic digest.
79. (newly added) The method of claim 28 wherein the label is a reactive label.
80. (newly added) A method for detecting one or more phosphopeptides or phosphoproteins in one or more samples containing a mixture of peptides or proteins which comprises the steps of:
- a. reacting the peptides or proteins in the mixture with a protective group that is an amine that reacts to protect any phosphate groups therein by forming phosphoramidate bonds and to protect any carboxylic acid groups and esters thereof therein by forming amide bonds;
 - b. treating the protected peptides or proteins in the mixture under conditions which selectively substantially cleave the phosphoramidate bonds, without substantially cleaving the amide bonds, to regenerate free phosphate groups in the peptide or protein;
 - c. selectively labeling the free phosphate groups in the peptides or proteins in the sample with a label having a functional group that reacts directly or indirectly with a phosphate group; and
 - d. detecting the peptides or proteins carrying the selective label of step c to detect the one or more phosphopeptides or phosphoproteins in the sample.

C₁

Ci
cont

81. (newly added) The method of claim 80 wherein the protective group is ethanol amine.
